Student Picture Here

EMERGENCY CARE PLAN - ALLE	KGY
Name Birth	date
School School Year	
Allergy to	
Asthma: ☐ Yes (higher risk for a severe reaction) ☐ N	No
ALL STUDENT'S EMERGENCY MEDICATIONS MUST BE EMERGENCY MEDICATIONS MUST ACCOMPANY STUDENT O	
Inject epinephrine immediately for an (with or without sym	
Circle specific symptoms listed below that your child is known to have: MILD SYMPTOMS ONLY: Exposure unknown MOUTH: Itchy mouth NOSE: Runny nose, sneezing SKIN: A few hives around mouth/face, mild itch in one area EYES: Red, watery, itching GUT: Mild nausea/discomfort	1. Administer antihistamine: Medication Location Dose When to give 2. Additional Medication if any: Medication Location Dose When to give 3. Stay with Student 4. Call parent
Circle specific symptoms listed below that your child is known to have: Any SEVERE SYMPTOMS: Exposure suspected or known One or more of the following: LUNG: Short of breath, wheeze, repetitive cough HEART: Pale, blue, faint, weak pulse, dizzy, confused THROAT: Tight, hoarse, trouble breathing/swallowing, drooling MOUTH: Obstructive swelling (tongue and/or lips) SKIN: Progressing hives, rash, swelling MENTAL: Feeling of doom, confusion, loss of consciousness, dizziness Or combination of symptoms for different body areas: SKIN: Hives, itchy rashes, swelling (eyes, lips) GUT: Vomiting, diarrhea, cramps	If symptoms don't improve or get worse, move on to Severe Symptom treatment 1. INJECT EPINEPHRINE IMMEDIATELY Medication Location Dose 2. Call 911, Notify time Epinephrine given. 3. Keep student calm and seated 4. Monitor students condition and provide first aid if necessary 5. If symptoms don't improve aftermin, Give second dose of EPI (if available) Location Medication Dose 6. Call parent 7. Stay with Student until EMS arrives 8. Complete District required forms
SKIN : Hives, itchy rashes, swelling (eyes, lips)	6. Call parent7. Stay with Student until EMS arrives

EPI PEN AND EPI PEN JR. ADMINISTRATION INSTRUCTIONS




- Hold firmly with orange tip pointing downward.
- Remove blue safety cap by pulling straight up. Do not bend or twist.




- Swing and push orange tip firmly into mid-outer thigh until you hear a "click."
- Hold on thigh for 10 seconds.


Auvi-Q™ Administration Guide



Pull cartridge from case.



Pull off **RED** Safety Guard



Place **BLACK** end against outer thigh, then press firmly and hold for 5 seconds.

Updated: 06.19.15